

Temporal Justification Logic

Samuel Bucheli

Zhlke Engineering AG,
Bogenschtzenstrasse 9A,
3008 Bern, Switzerland
samuel.bucheli.cs@gmail.com

Meghdad Ghari*

School of Mathematics,
Institute for Research
in Fundamental Sciences (IPM),
P.O. Box: 19395-5746, Tehran, Iran
ghari@ipm.ir

Thomas Studer†

Institut für Informatik,
Universität Bern,
Neubrückstrasse 10,
3012 Bern, Switzerland
tstuder@inf.unibe.ch

Justification logics are modal-like logics with the additional capability of recording the reason, or justification, for modalities in syntactic structures, called justification terms. Justification logics can be seen as explicit counterparts to modal logics. The behavior and interaction of agents in distributed system is often modeled using logics of knowledge and time. In this paper, we sketch some preliminary ideas on how the modal knowledge part of such logics of knowledge and time could be replaced with an appropriate justification logic.

1 Introduction

Justification logics are epistemic logics that feature explicit reasons for an agent's knowledge and belief. Originally, Artemov [1] developed the first justification logic, the Logic of Proofs LP, to provide a classical provability semantics for intuitionistic logic. Later, Fitting [11] introduced epistemic models for justification logic. This general reading of justification led to a big variety of epistemic justification logics for many different applications [2, 3, 4, 6, 13, 14, 19, 20]. Instead of an implicit statement $K\phi$, which stands for *the agent knows ϕ* , justification logics include explicit statements of the form $[t]\phi$, which mean *t justifies the agent's knowledge of ϕ* .

A common approach to model distributed systems of interacting agents is using logics of knowledge and time, with the interplay between these two modalities leading to interesting properties and questions [10, 17, 18, 21, 9]. While knowledge in such systems has typically been modeled using the modal logic S5, it is a natural question to ask what happens when we model knowledge in such logics using a justification logic.

This paper offers a first study on combining temporal logic and justification logic. We introduce a system $LPLTL_{CS}$ that combines linear time temporal logic LTL with the justification logic LP. In Sections 2 and 3 we present the language and the axioms of $LPLTL_{CS}$, respectively. In Section 4 we introduce interpreted systems with Fitting models as semantics for temporal justification logic. In Section 5 we establish soundness and completeness of $LPLTL_{CS}$. In Section 6 we present an extension $LPLTL_{CS}^*$ of $LPLTL_{CS}$ that enjoys the internalization property. In Section 7 we introduce some additional principles concerning interactions of knowledge, justifications, and time. In Section 8 we conclude the paper and discuss some open problems.

Acknowledgments. We would like to thank the anonymous referees for many helpful comments, which helped to improve the paper.

*This research was in part supported by a grant from IPM (No. 95030416).

†This work was partially supported by the SNSF project 200021_165549 *Justifications and non-classical reasoning*.

2 Language

In the following, let h be a fixed number of agents, Const a countable set of justification constants, Var a countable set of justification variables, and Prop a countable set of atomic propositions.

The set of justification terms Tm is defined inductively by

$$t ::= c \mid x \mid !t \mid t + t \mid t \cdot t,$$

where $c \in \text{Const}$ and $x \in \text{Var}$.

The set of formulas Fml is inductively defined by

$$\varphi ::= P \mid \perp \mid \varphi \rightarrow \varphi \mid \bigcirc \varphi \mid \varphi \mathcal{U} \varphi \mid [t]_i \varphi,$$

where $1 \leq i \leq h$, $t \in \text{Tm}$, and $P \in \text{Prop}$.

We use the following usual abbreviations:

$$\begin{aligned} \neg \varphi &:= \varphi \rightarrow \perp & \top &:= \neg \perp \\ \varphi \vee \psi &:= \neg \varphi \rightarrow \psi & \varphi \wedge \psi &:= \neg(\neg \varphi \vee \neg \psi) \\ \varphi \leftrightarrow \psi &:= (\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi) & \Diamond \varphi &:= \top \mathcal{U} \varphi \\ \Box \varphi &:= \neg \Diamond \neg \varphi. \end{aligned}$$

Associativity and precedence of connectives, as well as the corresponding omission of brackets, are handled in the usual manner.

Subformulas are defined as usual. The set of subformulas $\text{Sub}(\chi)$ of a formula χ is inductively given by:

$$\begin{aligned} \text{Sub}(P) &:= \{P\} & \text{Sub}(\perp) &:= \{\perp\} \\ \text{Sub}(\varphi \rightarrow \psi) &:= \{\varphi \rightarrow \psi\} \cup \text{Sub}(\varphi) \cup \text{Sub}(\psi) & \text{Sub}(\bigcirc \varphi) &:= \{\bigcirc \varphi\} \cup \text{Sub}(\varphi) \\ \text{Sub}(\varphi \mathcal{U} \psi) &:= \{\varphi \mathcal{U} \psi\} \cup \text{Sub}(\varphi) \cup \text{Sub}(\psi) & \text{Sub}([t]_i \varphi) &:= \{[t]_i \varphi\} \cup \text{Sub}(\varphi). \end{aligned}$$

3 Axioms

The axiom system for temporal justification logic consists of three parts, namely propositional logic, temporal logic, and justification logic.

Propositional Logic

For propositional logic, we take

1. all propositional tautologies (Taut)

as axioms and the rule modus ponens, as usual:

$$\frac{\varphi \quad \varphi \rightarrow \psi}{\psi} (\text{MP}).$$

Temporal Logic

For the temporal part, we use a system of [12, 15, 16] with axioms

2. $\bigcirc(\varphi \rightarrow \psi) \rightarrow (\bigcirc\varphi \rightarrow \bigcirc\psi)$ (\bigcirc -k)
3. $\Box(\varphi \rightarrow \psi) \rightarrow (\Box\varphi \rightarrow \Box\psi)$ (\Box -k)
4. $\bigcirc\neg\varphi \leftrightarrow \neg\bigcirc\varphi$ (fun)
5. $\Box(\varphi \rightarrow \bigcirc\varphi) \rightarrow (\varphi \rightarrow \Box\varphi)$ (ind)
6. $\varphi \mathcal{U} \psi \rightarrow \Diamond\psi$ (\mathcal{U} 1)
7. $\varphi \mathcal{U} \psi \leftrightarrow \psi \vee (\varphi \wedge \bigcirc(\varphi \mathcal{U} \psi))$ (\mathcal{U} 2)

and rules

$$\frac{\varphi}{\bigcirc\varphi} (\bigcirc\text{-nec}), \quad \frac{\varphi}{\Box\varphi} (\Box\text{-nec}).$$

Justification Logic

Finally, for the justification logic part, we use a multi-agent version of the Logic of Proofs [1, 6, 13, 27] with axioms

8. $[t]_i(\varphi \rightarrow \psi) \rightarrow ([s]_i\varphi \rightarrow [t \cdot s]_i\psi)$ (application)
9. $[t]_i\varphi \rightarrow [t + s]_i\varphi \quad [s]_i\varphi \rightarrow [t + s]_i\varphi$ (sum)
10. $[t]_i\varphi \rightarrow \varphi$ (reflexivity)
11. $[t]_i\varphi \rightarrow [!t]_i[t]_i\varphi$ (positive introspection)

and rule

$$\frac{[c]_i\varphi \in \text{CS}}{[c]_i\varphi} (\text{ax-nec}),$$

where the constant specification CS is a set of formulas $[c]_i\varphi$, where $c \in \text{Const}$ is a justification constant and φ is an axiom of propositional logic, temporal logic, or justification logic.

For a given constant specification CS, we use LPLTL_{CS} to denote the Hilbert system given by the axioms and rules for propositional logic, temporal logic, and justification logic as presented above. As usual, we write $\text{LPLTL}_{\text{CS}} \vdash \varphi$ or simply $\vdash_{\text{CS}} \varphi$ if a formula φ is derivable in LPLTL_{CS} . Often the constant specification is clear from the context and we will only write $\vdash \varphi$ instead of $\vdash_{\text{CS}} \varphi$.

The axiomatization for linear time temporal logic given in [12, 15, 16] includes an axiom

$$\Box\varphi \rightarrow (\varphi \wedge \bigcirc\Box\varphi).$$

The following lemma shows that we do not need this axiom since in our formalization \Box is a defined operator.

Lemma 1. *We have*

$$\vdash_{\text{CS}} \Box\varphi \rightarrow (\varphi \wedge \bigcirc\Box\varphi)$$

and (MP) is the only rule that is used in this derivation.

Proof. $\Box\varphi$ stands for $\neg(\top \mathcal{U} \neg\varphi)$. Hence from ($\mathcal{U}2$) we get

$$\vdash_{CS} \neg\varphi \vee \bigcirc(\top \mathcal{U} \neg\varphi) \rightarrow \top \mathcal{U} \neg\varphi.$$

Taking the contrapositive yields

$$\vdash_{CS} \neg(\top \mathcal{U} \neg\varphi) \rightarrow \neg(\neg\varphi \vee \bigcirc(\top \mathcal{U} \neg\varphi)).$$

By propositional reasoning and (fun) we get

$$\vdash_{CS} \neg(\top \mathcal{U} \neg\varphi) \rightarrow (\varphi \wedge \bigcirc\neg(\top \mathcal{U} \neg\varphi)),$$

which is

$$\vdash_{CS} \Box\varphi \rightarrow (\varphi \wedge \bigcirc\Box\varphi). \quad \square$$

Remark. As usual, we find that the following rule is derivable, see [5, Lemma 6] for a detailed derivation,

$$\frac{\chi \rightarrow \neg\psi \wedge \bigcirc\chi}{\chi \rightarrow \neg(\varphi \mathcal{U} \psi)}.$$

From this, we get that the following rule is also derivable

$$\frac{\chi \rightarrow \neg\psi \wedge \bigcirc(\chi \vee (\neg\varphi \wedge \neg\psi))}{\chi \rightarrow \neg(\varphi \mathcal{U} \psi)} (\mathcal{U}-R).$$

A proof is given in [17, Lemma 4.5].

4 Semantics

In this section we introduce interpreted systems based on Fitting-models as semantics for temporal justification logic.

Definition 2. A frame is a tuple (S, R_1, \dots, R_h) where

1. S is a non-empty set of states;
2. each $R_i \subseteq S \times S$ is a reflexive and transitive relation.

A run r on a frame is a function from \mathbb{N} to states, i.e., $r : \mathbb{N} \rightarrow S$. A system \mathcal{R} is a non-empty set of runs.

Definition 3. Given a frame (S, R_1, \dots, R_h) , a CS-evidence function for agent i is a function

$$\mathcal{E}_i : S \times Tm \rightarrow \mathcal{P}(Fml)$$

satisfying the following conditions. For all terms $s, t \in Tm$, all formulas $\varphi, \psi \in Fml$, and all $v, w \in S$,

1. $\mathcal{E}_i(v, t) \subseteq \mathcal{E}_i(w, t)$, whenever $R_i(v, w)$ (monotonicity)
2. if $[c]_i \varphi \in CS$, then $\varphi \in \mathcal{E}_i(w, c)$ (constant specification)
3. if $\varphi \rightarrow \psi \in \mathcal{E}_i(w, t)$ and $\varphi \in \mathcal{E}_i(w, s)$, then $\psi \in \mathcal{E}_i(w, t \cdot s)$ (application)
4. $\mathcal{E}_i(w, s) \cup \mathcal{E}_i(w, t) \subseteq \mathcal{E}_i(w, s + t)$ (sum)
5. if $\varphi \in \mathcal{E}_i(w, t)$, then $[t]_i \varphi \in \mathcal{E}_i(w, !t)$ (positive introspection)

Definition 4. An interpreted system for CS is a tuple

$$\mathcal{I} = (\mathcal{R}, S, R_1, \dots, R_h, \mathcal{E}_1, \dots, \mathcal{E}_h, v)$$

where

1. (S, R_1, \dots, R_h) is a frame;
2. \mathcal{R} is a system on that frame;
3. \mathcal{E}_i is a CS-evidence function for agent i for $1 \leq i \leq h$;
4. $v : S \rightarrow \mathcal{P}(\text{Prop})$ is a valuation.

Definition 5. Given an interpreted system

$$\mathcal{I} = (\mathcal{R}, S, R_1, \dots, R_h, \mathcal{E}_1, \dots, \mathcal{E}_h, v),$$

a run $r \in \mathcal{R}$, and $n \in \mathbb{N}$, we define truth of a formula ϕ in \mathcal{I} at state $r(n)$ inductively by

$$\begin{aligned} (\mathcal{I}, r, n) &\models P \text{ iff } P \in v(r(n)), \\ (\mathcal{I}, r, n) &\not\models \perp, \\ (\mathcal{I}, r, n) &\models \phi \rightarrow \psi \text{ iff } (\mathcal{I}, r, n) \not\models \phi \text{ or } (\mathcal{I}, r, n) \models \psi, \\ (\mathcal{I}, r, n) &\models \bigcirc \phi \text{ iff } (\mathcal{I}, r, n+1) \models \phi, \\ (\mathcal{I}, r, n) &\models \phi \mathcal{U} \psi \text{ iff there is some } m \geq 0 \text{ such that } (\mathcal{I}, r, n+m) \models \psi \\ &\quad \text{and } (\mathcal{I}, r, n+k) \models \phi \text{ for all } 0 \leq k < m, \\ (\mathcal{I}, r, n) &= [t]_i \phi \text{ iff } \phi \in \mathcal{E}_i(r(n), t) \text{ and } (\mathcal{I}, r', n') \models \phi \\ &\quad \text{for all } r' \in \mathcal{R} \text{ and } n' \in \mathbb{N} \text{ such that } R_i(r(n), r'(n')). \end{aligned}$$

As usual, we write $\mathcal{I} \models \phi$ if for all $r \in \mathcal{R}$ and all $n \in \mathbb{N}$, we have $(\mathcal{I}, r, n) \models \phi$. Further, we write $\models_{CS} \phi$ if $\mathcal{I} \models \phi$ for all interpreted systems \mathcal{I} for CS.

Remark. From the definitions of \Box and \Diamond it follows that:

$$\begin{aligned} (\mathcal{I}, r, n) &\models \Diamond \phi \text{ iff } (\mathcal{I}, r, n+k) \models \phi \text{ for some } k \geq 0, \\ (\mathcal{I}, r, n) &\models \Box \phi \text{ iff } (\mathcal{I}, r, n+k) \models \phi \text{ for all } k \geq 0. \end{aligned}$$

5 Soundness and Completeness

The soundness proof for LPLTL_{CS} is a straightforward combination of the soundness proofs for temporal logic and justification logic by induction on the derivation.

Theorem 6. Let CS be an arbitrary constant specification. For each formula ϕ ,

$$\vdash_{CS} \phi \text{ implies } \models_{CS} \phi.$$

Our completeness proof for LPLTL_{CS} follows the one given in [17]. First, we define

$$\Gamma \vdash_{CS} \phi \text{ iff there exist } \psi_1, \dots, \psi_n \in \Gamma \text{ such that } \vdash_{CS} (\psi_1 \wedge \dots \wedge \psi_n) \rightarrow \phi.$$

Following our convention, we will usually write $\Gamma \vdash \phi$ instead of $\Gamma \vdash_{CS} \phi$.

Definition 7. Let CS be a constant specification. A set Γ of formulas is called CS -consistent if $\Gamma \not\vdash_{CS} \perp$. That means $\not\vdash_{CS} \bigwedge \Sigma \rightarrow \perp$, for each finite $\Sigma \subseteq \Gamma$.

For a formula χ , let $\text{Sub}^+(\chi) := \text{Sub}(\chi) \cup \{\neg\psi \mid \psi \in \text{Sub}(\chi)\}$. Let MCS_χ denote the set of all maximally CS -consistent subsets of $\text{Sub}^+(\chi)$. We have the following facts for $\Gamma \in \text{MCS}_\chi$:

- If $\Gamma \vdash_{CS} \varphi$, then $\vdash_{CS} \bigwedge \Gamma \rightarrow \varphi$.
- If $\varphi \in \text{Sub}(\chi)$ and $\varphi \notin \Gamma$, then $\neg\varphi \in \Gamma$.
- If $\varphi \in \text{Sub}^+(\chi)$ and $\Gamma \vdash_{CS} \varphi$, then $\varphi \in \Gamma$.
- If $\psi \in \text{Sub}^+(\chi)$, $\varphi \in \Gamma$ and $\vdash_{CS} \varphi \rightarrow \psi$, then $\psi \in \Gamma$.

We define the relation R_\circ on MCS_χ as follows:

$$\Gamma R_\circ \Delta \quad \text{iff} \quad \not\vdash_{CS} \bigwedge \Gamma \rightarrow \neg \bigcirc \bigwedge \Delta.$$

From this definition we immediately get the following lemmas.

Lemma 8. Let $\Gamma, \Delta \in \text{MCS}_\chi$, $\Gamma R_\circ \Delta$, and $\varphi \in \text{Sub}(\chi)$.

1. If $\Gamma \vdash_{CS} \bigcirc \varphi$, then $\varphi \in \Delta$.
2. If $\Gamma \vdash_{CS} \neg \bigcirc \varphi$, then $\neg \varphi \in \Delta$.

Proof. 1. Suppose toward a contradiction that $\varphi \notin \Delta$. Thus $\neg\varphi \in \Delta$. Since $\Gamma \vdash_{CS} \bigcirc \varphi$, we have $\vdash_{CS} \bigwedge \Gamma \rightarrow \bigcirc \varphi$. Hence $\vdash_{CS} \bigwedge \Gamma \rightarrow \bigcirc \neg \neg \varphi$. Therefore $\vdash_{CS} \bigwedge \Gamma \rightarrow \bigcirc \neg \bigwedge \Delta$. Thus

$$\vdash_{CS} \bigwedge \Gamma \rightarrow \neg \bigcirc \bigwedge \Delta,$$

which would contradict $\Gamma R_\circ \Delta$.

2. The proof is similar to part 1. □

Lemma 9. Let $\Gamma \in \text{MCS}_\chi$ and let $S := \{\Delta \in \text{MCS}_\chi \mid \Gamma R_\circ \Delta\}$. We have

$$\vdash \bigwedge \Gamma \rightarrow \bigcirc \bigvee \{ \bigwedge \Delta \mid \Delta \in S \}.$$

Proof. First observe that for all $\Gamma, \Delta \in \text{MCS}_\chi$ we have

$$(\text{not } \Gamma R_\circ \Delta) \quad \text{implies} \quad \vdash \bigwedge \Gamma \rightarrow \neg \bigcirc \bigwedge \Delta. \quad (1)$$

We also have

$$\vdash \bigvee \{ \bigwedge \Delta \mid \Delta \in \text{MCS}_\chi \}.$$

By necessitation we get

$$\vdash \bigcirc \bigvee \{ \bigwedge \Delta \mid \Delta \in \text{MCS}_\chi \}$$

and thus

$$\vdash \bigvee \{ \bigcirc \bigwedge \Delta \mid \Delta \in \text{MCS}_\chi \}. \quad (2)$$

By (1) we infer

$$\vdash \bigwedge \Gamma \rightarrow \bigvee \{ \bigcirc \bigwedge \Delta \mid \Delta \in \text{MCS}_\chi \text{ with } \Gamma R_\circ \Delta \}$$

and thus

$$\vdash \bigwedge \Gamma \rightarrow \bigcirc \bigvee \{ \bigwedge \Delta \mid \Delta \in \text{MCS}_\chi \text{ with } \Gamma R_\circ \Delta \}. \quad \square$$

Lemma 10. *The relation R_\circ is serial. That is for each $\Gamma \in \text{MCS}_\chi$, there exists $\Delta \in \text{MCS}_\chi$ with $\Gamma R_\circ \Delta$.*

Proof. Suppose towards a contradiction that for $\Gamma \in \text{MCS}_\chi$ we have $(\text{not } \Gamma R_\circ \Delta)$ for all $\Delta \in \text{MCS}_\chi$. Then $\vdash \bigwedge \Gamma \rightarrow \neg \bigcirc \bigwedge \Delta$, for all $\Delta \in \text{MCS}_\chi$. Thus

$$\vdash \bigwedge \Gamma \rightarrow \bigwedge \{\neg \bigcirc \bigwedge \Delta \mid \Delta \in \text{MCS}_\chi\},$$

and hence,

$$\vdash \bigwedge \Gamma \rightarrow \neg \bigvee \{\bigcirc \bigwedge \Delta \mid \Delta \in \text{MCS}_\chi\}. \quad (3)$$

On the other hand, from (2) we deduce

$$\vdash \bigwedge \Gamma \rightarrow \bigvee \{\bigcirc \bigwedge \Delta \mid \Delta \in \text{MCS}_\chi\}. \quad (4)$$

Since Γ is consistent, (3) and (4) leads to a contradiction. \square

Definition 11. *A finite sequence $(\Gamma_0, \Gamma_1, \dots, \Gamma_n)$ of elements of MCS_χ is called a $\varphi \mathcal{U} \psi$ -sequence starting with Γ if*

1. $\Gamma_0 = \Gamma$,
2. $\Gamma_j R_\circ \Gamma_{j+1}$, for all $j < n$,
3. $\psi \in \Gamma_n$,
4. $\varphi \in \Gamma_j$, for all $j < n$.

Lemma 12. *For every $\Gamma \in \text{MCS}_\chi$, if $\varphi \mathcal{U} \psi \in \Gamma$, then there exists a $\varphi \mathcal{U} \psi$ -sequence starting with Γ .*

Proof. Suppose $\varphi \mathcal{U} \psi \in \Gamma$ and there exists no $\varphi \mathcal{U} \psi$ -sequence starting with Γ . We let T be the smallest set of elements of MCS_χ such that

1. $\Gamma \in T$;
2. for each $\Delta' \in \text{MCS}_\chi$, if $\Delta \in T$, $\Delta R_\circ \Delta'$, and $\varphi \in \Delta'$, then $\Delta' \in T$.

We find that $\vdash \bigwedge \Delta \rightarrow \neg \psi$ for all $\Delta \in T$. Let

$$\rho := \bigvee \{\bigwedge \Delta \mid \Delta \in T\}.$$

We have $\vdash \rho \rightarrow \neg \psi$.

Moreover, for each $\Delta \in T$ and each $\Delta' \in \text{MCS}_\chi$ with $\Delta R_\circ \Delta'$, we have

$$\text{either } \Delta' \in T \text{ or } \vdash \bigwedge \Delta' \rightarrow \neg \varphi \wedge \neg \psi.$$

Thus, by Lemma 9, we get $\vdash \rho \rightarrow \bigcirc(\rho \vee (\neg \varphi \wedge \neg \psi))$. Using (\mathcal{U} -R), we obtain $\vdash \rho \rightarrow \neg(\varphi \mathcal{U} \psi)$. Since $\Gamma \in T$, this implies $\vdash \bigwedge \Gamma \rightarrow \neg(\varphi \mathcal{U} \psi)$, which contradicts the assumption $\varphi \mathcal{U} \psi \in \Gamma$. \square

Definition 13. *An infinite sequence $(\Gamma_0, \Gamma_1, \dots)$ of elements of MCS_χ is called acceptable if*

1. $\Gamma_n R_\circ \Gamma_{n+1}$ for all $n \geq 0$, and
2. for all n , if $\varphi \mathcal{U} \psi \in \Gamma_n$, then there exists $m \geq n$ such that $\psi \in \Gamma_m$ and $\varphi \in \Gamma_k$ for all k with $n \leq k < m$.

Lemma 14. *Every finite sequence $(\Gamma_0, \Gamma_1, \dots, \Gamma_n)$ of elements of MCS_χ with $\Gamma_j R_\circ \Gamma_{j+1}$, for all $j < n$, can be extended to an acceptable sequence.*

Proof. In order to fulfill the requirements of Definition 13, we shall extend the sequence $(\Gamma_0, \Gamma_1, \dots, \Gamma_n)$ by the following algorithm.

Suppose $\varphi \mathcal{U} \psi \in \Gamma_0$. Then either $\psi \in \Gamma_0$ or $\neg\psi \in \Gamma_0$. In the former case the requirement is fulfilled for the formula $\varphi \mathcal{U} \psi$ in Γ_0 , and we go to the next step. In the latter case, using axiom ($\mathcal{U} 2$),

$$\Gamma_0 \vdash_{CS} \varphi \wedge \bigcirc(\varphi \mathcal{U} \psi).$$

Since $\Gamma_0 R_{\bigcirc} \Gamma_1$, by Lemma 8, we get $\varphi \mathcal{U} \psi \in \Gamma_1$.

We can repeat this argument for Γ_i for $1 \leq i \leq n$. We find that the requirement for $\varphi \mathcal{U} \psi \in \Gamma_0$ is either fulfilled in $(\Gamma_0, \Gamma_1, \dots, \Gamma_n)$ or $\varphi \mathcal{U} \psi \in \Gamma_n$ and $\varphi \in \Gamma_i$ for $1 \leq i \leq n$. In the latter case, by Lemma 12, there exists a sequence $(\Gamma_n, \Gamma_{n+1}, \dots, \Gamma_{n+m})$ such that $\varphi \in \Gamma_i$ for $n \leq i < n+m$, $\psi \in \Gamma_{n+m}$, and $\Gamma_i R_{\bigcirc} \Gamma_{i+1}$ for $n \leq i < n+m$. This gives a finite extension of the original sequence that satisfies the requirement imposed by $\varphi \mathcal{U} \psi \in \Gamma_0$.

In the next step we repeat this argument for the remaining obligations at Γ_0 . Eventually we obtain a finite sequence that satisfies all requirements imposed by formulas at Γ_0 .

We may move on to Γ_1 and apply the same procedure. It is clear that by iterating it we obtain in the limit an acceptable sequence that extends $(\Gamma_0, \Gamma_1, \dots, \Gamma_n)$. \square

Corollary 15. *For every $\Gamma \in \text{MCS}_\chi$, there is an acceptable sequence that starts with Γ .*

Definition 16. *The χ -canonical interpreted system*

$$\mathcal{I} = (\mathcal{R}, S, R_1, \dots, R_h, \mathcal{E}_1, \dots, \mathcal{E}_h, \nu)$$

for CS is defined as follows:

1. \mathcal{R} consists of all mappings $r : \mathbb{N} \rightarrow \text{MCS}_\chi$ such that $(r(0), r(1), \dots)$ is an acceptable sequence;
2. $S := \text{MCS}_\chi = \{r(n) \mid r \in \mathcal{R}, n \in \mathbb{N}\}$;
3. $R_i(\Gamma, \Delta)$ iff $\{\varphi \mid \Gamma \vdash [t]_i \varphi \text{ for some } t\} \subseteq \{\varphi \mid \Delta \vdash \varphi\}$;
4. $\mathcal{E}_i(\Gamma, t) := \{\varphi \mid \Gamma \vdash [t]_i \varphi\}$;
5. $\nu(\Gamma) := \{P \in \text{Prop} \mid P \in \Gamma\}$.

Remark. *The χ -canonical interpreted system \mathcal{I} for CS is a finite structure in the sense that the set of states S is finite. This is a novelty for completeness proofs of justification logics. Even the completeness proofs for justification logics with common knowledge [2, 6] work with infinite canonical structures. Note that this remark concerns epistemic Fitting-models. Of course, symbolic M-models [22] could be considered as single-world Fitting-models.*

The fact that states of \mathcal{I} are maximally CS-consistent subsets of $\text{Sub}^+(\chi)$ —instead of just maximally CS-consistent sets—matters for the definitions of R_i and \mathcal{E}_i . The usual definitions would be

$$R_i(\Gamma, \Delta) \text{ iff } \{\varphi \mid [t]_i \varphi \in \Gamma \text{ for some } t\} \subseteq \{\varphi \mid \varphi \in \Delta\} \quad \text{and} \\ \mathcal{E}_i(\Gamma, t) := \{\varphi \mid [t]_i \varphi \in \Gamma\}.$$

This, however, would not work for our finite canonical structure. In particular the next lemma could not be established as, for instance, $[t]_i \varphi \in \Gamma$ does not imply $[!t]_i [t]_i \varphi \in \Gamma$ for $\Gamma \in \text{MCS}_\chi$.

Lemma 17. *The χ -canonical interpreted system*

$$\mathcal{I} = (\mathcal{R}, S, R_1, \dots, R_h, \mathcal{E}_1, \dots, \mathcal{E}_h, \nu)$$

for CS is an interpreted system for CS.

Proof. The proof is essentially the same as the corresponding proof for single agent Fitting-models in [11]. Let us only show here the monotonicity condition for \mathcal{E}_i .

Suppose $\Gamma, \Delta \in S$ and $R_i(\Gamma, \Delta)$. Suppose that $\varphi \in \mathcal{E}_i(\Gamma, t)$. Thus $\Gamma \vdash [t]_i \varphi$. Hence $\Gamma \vdash [!t]_i [t]_i \varphi$. Since $R_i(\Gamma, \Delta)$, we have $\Delta \vdash [t]_i \varphi$. Hence $\varphi \in \mathcal{E}_i(\Delta, t)$ as desired. \square

Lemma 18 (Truth Lemma). *Let $\mathcal{J} = (\mathcal{R}, S, R_1, \dots, R_h, \mathcal{E}_1, \dots, \mathcal{E}_h, \nu)$ be the χ -canonical interpreted system for CS. For every formula $\psi \in \text{Sub}^+(\chi)$, every run r in \mathcal{R} , and every $n \in \mathbb{N}$ we have:*

$$(\mathcal{J}, r, n) \models \psi \quad \text{iff} \quad \psi \in r(n).$$

Proof. As usual, the proof is by induction on the structure of ψ . We show only the following cases:

- $\psi = [t]_i \varphi$. (\Rightarrow) If $(\mathcal{J}, r, n) \models [t]_i \varphi$, then $\varphi \in \mathcal{E}_i(r(n), t)$. Thus, by definition, $r(n) \vdash [t]_i \varphi$. Hence $[t]_i \varphi \in r(n)$, since $[t]_i \varphi \in \text{Sub}^+(\chi)$.
 (\Leftarrow) If $[t]_i \varphi \in r(n)$, then $r(n) \vdash [t]_i \varphi$. Hence, by definition, $\varphi \in \mathcal{E}_i(r(n), t)$. Now suppose that $R_i(r(n), r'(n'))$. We find $r'(n') \vdash \varphi$. Since $\varphi \in \text{Sub}^+(\chi)$, we have $\varphi \in r'(n')$ and by I.H. we get $(\mathcal{J}, r', n') \models \varphi$. Since r' and n' were arbitrary, we conclude $(\mathcal{J}, r, n) \models [t]_i \varphi$.
- $\psi = \bigcirc \varphi$. (\Rightarrow) Suppose that $(\mathcal{J}, r, n) \models \bigcirc \varphi$ and $\bigcirc \varphi \notin r(n)$. Then $(\mathcal{J}, r, n+1) \models \varphi$, and hence by the induction hypothesis $\varphi \in r(n+1)$. On the other hand, $\neg \bigcirc \varphi \in r(n)$. Since $r(n) R_{\bigcirc} r(n+1)$, by Lemma 8, we get $\neg \varphi \in r(n+1)$, which is a contradiction.
 (\Leftarrow) If $\bigcirc \varphi \in r(n)$, then $\varphi \in r(n+1)$. By the induction hypothesis, $(\mathcal{J}, r, n+1) \models \varphi$, and hence $(\mathcal{J}, r, n) \models \bigcirc \varphi$.
- $\psi = \psi_1 \mathcal{U} \psi_2$. (\Rightarrow) If $(\mathcal{J}, r, n) \models \psi_1 \mathcal{U} \psi_2$, then $(\mathcal{J}, r, m) \models \psi_2$ for some $m \geq n$, and $(\mathcal{J}, r, k) \models \psi_1$ for all k with $n \leq k < m$. By I.H. we get $\psi_2 \in r(m)$, and $\psi_1 \in r(k)$ for all k with $n \leq k < m$. We have to show $\psi_1 \mathcal{U} \psi_2 \in r(n)$, which follows by induction on m as follows:
 - Base case $m = n$. Since $\psi_2 \in r(n)$ and $\vdash \psi_2 \rightarrow (\psi_1 \mathcal{U} \psi_2)$, we obtain $\psi_1 \mathcal{U} \psi_2 \in r(n)$.
 - Suppose $m > n$. It follows from the induction hypothesis that $\psi_1 \mathcal{U} \psi_2 \in r(n+1)$. From this and $r(n) R_{\bigcirc} r(n+1)$ we get that

$$r(n) \cup \{\bigcirc(\psi_1 \mathcal{U} \psi_2)\} \text{ is consistent.} \quad (5)$$

Assume now

$$\neg(\psi_1 \mathcal{U} \psi_2) \in r(n). \quad (6)$$

Then $r(n) \vdash \neg(\psi_1 \mathcal{U} \psi_2)$ and by axiom ($\mathcal{U} 2$) we find $r(n) \vdash \neg(\psi_1 \wedge \bigcirc(\psi_1 \mathcal{U} \psi_2))$. From $\psi_1 \in r(n)$ we get $r(n) \vdash \psi_1$ and thus $r(n) \vdash \neg \bigcirc(\psi_1 \mathcal{U} \psi_2)$, which contradicts (5). Hence the assumption (6) must be false and we conclude $\psi_1 \mathcal{U} \psi_2 \in r(n)$.

(\Leftarrow) If $\psi_1 \mathcal{U} \psi_2 \in r(n)$, then since $(r(n), r(n+1), \dots)$ is an acceptable sequence there exists $m \geq n$ such that $\psi_2 \in r(m)$, and $\psi_1 \in r(k)$ for all k with $n \leq k < m$. By I.H. we obtain $(\mathcal{J}, r, m) \models \psi_2$, and $(\mathcal{J}, r, k) \models \psi_1$ for all k with $n \leq k < m$. Thus $(\mathcal{J}, r, n) \models \psi_1 \mathcal{U} \psi_2$. \square

Theorem 19 (Completeness). *For each formula φ ,*

$$\models_{\text{CS}} \varphi \quad \text{implies} \quad \vdash_{\text{CS}} \varphi.$$

Proof. Suppose that $\not\models_{\text{CS}} \varphi$. Thus, $\{\neg \varphi\}$ is a CS-consistent set. Therefore, there exists $\Gamma \in \text{MCS}_{\varphi}$ with $\neg \varphi \in \Gamma$. By Corollary 15, there is an acceptable sequence starting with Γ . Thus there is a run r in the φ -canonical interpreted system \mathcal{J} for CS with $r(0) = \Gamma$. Since $\neg \varphi \in \Gamma$, by the Truth Lemma, $(\mathcal{J}, r, 0) \not\models \varphi$. Therefore, $\not\models_{\text{CS}} \varphi$. \square

6 Internalization

It is desirable that a justification logic internalizes its own notion of proof. This is formalized in the following definition.

Definition 20. A justification logic L satisfies internalization if for each formula φ with $L \vdash \varphi$ and for each agent i , there exists a term t with $L \vdash [t]_i \varphi$.

Usually, internalization is shown by induction on the derivation of φ . However, for $LPLTL_{CS}$ this seems not possible because it includes rules (\bigcirc -nec) and (\Box -nec). In this section, we introduce an extension $LPLTL_{CS}^*$ of $LPLTL_{CS}$ that satisfies internalization.

The language of $LPLTL_{CS}^*$ includes a new unary operator \star on justification terms. We define

$$\star^0 c := c \quad \text{and} \quad \star^n c := \star \star^{n-1} c \quad (\text{for } n \geq 1).$$

The set of terms Tm^* of $LPLTL_{CS}^*$ is given by

$$t ::= \star^n c \mid x \mid !t \mid t + t \mid t \cdot t,$$

where $c \in \text{Const}$, $n \geq 0$, and $x \in \text{Var}$. The set of formulas Fml^* of $LPLTL_{CS}^*$ is defined like Fml but using Tm^* instead of Tm .

The axioms of $LPLTL_{CS}^*$ are:

1. all axioms of $LPLTL$
2. $\Box \varphi \rightarrow \bigcirc \varphi$ (mix)
3. $\Box([t]_i \varphi \rightarrow \varphi)$ (boxed reflexivity)

The rules of $LPLTL_{CS}^*$ are:

$$\frac{\varphi \quad \varphi \rightarrow \psi}{\psi} \text{ (MP)} \quad \text{and} \quad \frac{[c]_{i_0} \varphi \in CS}{[\star^n c]_{i_n} \Box [\star^{n-1} c]_{i_{n-1}} \Box \dots [\star c]_{i_1} \Box [c]_{i_0} \varphi} \text{ (ax-nec)}^*,$$

where $n \geq 0$; so $(\text{ax-nec})^*$ subsumes (ax-nec) . Note that a constant specification for $LPLTL^*$ may include formulas of the form $[c]_i(\Box \varphi \rightarrow \bigcirc \varphi)$ and $[c]_i \Box([t]_i \varphi \rightarrow \varphi)$.

Remark. The principles (mix) and (boxed reflexivity) are derivable in $LPLTL_{CS}$. However, their proofs require applications of the rules (\bigcirc -nec) and (\Box -nec), respectively. Since these rules are not included in $LPLTL_{CS}^*$, we have to include (mix) and (boxed reflexivity) as axioms.

Remark. The \star -operation is very powerful. Its meaning can be explained as follows. If $[c]_i \varphi$ is contained in CS , then $[c]_i \varphi$ is provable and hence $\Box [c]_i \varphi$ is provable, too (see Lemma 22). The evidence $\star c$ justifies this fact, i.e., $[\star c]_i \Box [c]_i \varphi$ is provable. Looking closely at $(\text{ax-nec})^*$ we see that we get even more. Indeed, for any agent j we have that $[\star c]_j \Box [c]_i \varphi$ is provable. Moreover, even arbitrary iterations of this principle are provable, which implies that the constant specification is common knowledge among the agents, so to speak.

We could use a less general version of $(\text{ax-nec})^*$ where the \star -operation is indexed. This would be similar to the evidence verification operation of [27], see also Question 2. In that case we would obtain $[\star_i^j c]_j \Box [c]_i \varphi$. However, for the purpose of internalization we do not need these indices and hence we dispense with them.

Definition 21. A constant specification CS is axiomatically appropriate if for each axiom φ of $LPLTL^*$ and each agent i , there is a constant c with $[c]_i \varphi \in CS$.

First we show that $LPLTL_{CS}^*$ extends $LPLTL_{CS}$.

Lemma 22. *Let CS be an axiomatically appropriate constant specification for $LPLTL^*$. The rules $(\Box\text{-nec})$ and $(\bigcirc\text{-nec})$ are derivable in $LPLTL_{CS}^*$.*

Proof. We first show that $(\Box\text{-nec})$ is derivable in $LPLTL_{CS}^*$. Suppose φ is provable in $LPLTL_{CS}^*$. By induction on the proof of φ , we show that $\Box\varphi$ is provable in $LPLTL_{CS}^*$.

In case φ is an axiom, since CS is axiomatically appropriate, there is a constant c such that $[c]_i\varphi \in CS$. Using $(\text{ax-nec})^*$, we get $[\star c]_i\Box[c]_i\varphi$, and then using axiom (reflexivity) we get $\Box[c]_i\varphi$. Finally, using axioms (boxed reflexivity) and $(\Box\text{-k})$ we obtain $\Box\varphi$.

In case φ is derived by modus ponens, the claim is immediate by $(\Box\text{-k})$.

In case φ is $[\star^n c]_{i_n}\Box[\star^{n-1}c]_{i_{n-1}}\Box\ldots[\star c]_{i_1}\Box[c]_{i_0}\varphi$ derived using $(\text{ax-nec})^*$, we can use $(\text{ax-nec})^*$ also to obtain

$$[\star^{n+1}c]_{i_{n+1}}\Box[\star^n c]_{i_n}\Box[\star^{n-1}c]_{i_{n-1}}\Box\ldots[\star c]_{i_1}\Box[c]_{i_0}\varphi.$$

Then using (reflexivity) we get

$$\Box[\star^n c]_{i_n}\Box[\star^{n-1}c]_{i_{n-1}}\Box\ldots[\star c]_{i_1}\Box[c]_{i_0}\varphi,$$

that is $\Box\varphi$.

Derivability of $(\bigcirc\text{-nec})$ follows from $(\Box\text{-nec})$ and axiom (mix). □

Let CS be a constant specification for $LPLTL^*$. We set

$$CS^r := \{[c]_i\varphi \mid [c]_i\varphi \in CS \text{ and } \varphi \text{ is an axiom of } LPLTL\}.$$

Obviously, CS^r is a constant specification for $LPLTL$. We get the following corollary.

Corollary 23. *Let CS be an axiomatically appropriate constant specification for $LPLTL^*$. For each formula φ of Fml ,*

$$LPLTL_{CS^r} \vdash \varphi \text{ implies } LPLTL_{CS}^* \vdash \varphi.$$

We will now establish the internalization property. We need the following lemma.

Lemma 24. *Let CS be an axiomatically appropriate constant specification. For each formula φ and each i ,*

$$LPLTL_{\emptyset}^* \vdash \varphi \text{ implies } LPLTL_{CS}^* \vdash [t]_i\varphi \text{ for some term } t.$$

Proof. We proceed by induction on the derivation of φ .

In case φ is an axiom, since CS is axiomatically appropriate, there is a constant c with

$$LPLTL_{CS}^* \vdash [c]_i\varphi.$$

In case φ is derived by modus ponens from $\psi \rightarrow \varphi$ and ψ , then, by the induction hypothesis, there are term s_1 and s_2 such that $[s_1]_i(\psi \rightarrow \varphi)$ and $[s_2]_i\psi$ are provable. Using (application) and modus ponens, we obtain $[s_1 \cdot s_2]_i\varphi$. □

Theorem 25. *Let CS be an axiomatically appropriate constant specification. $LPLTL_{CS}^*$ enjoys internalization.*

Proof. We have to show that for each formula φ and each i

$$\text{LPLTL}_{\text{CS}}^* \vdash \varphi \text{ implies } \text{LPLTL}_{\text{CS}}^* \vdash [t]_i \varphi \text{ for some term } t.$$

We proceed by induction on the derivation of φ .

The cases where φ is an axiom or φ is derived by modus ponens are like the corresponding cases in the previous lemma.

In case φ is $[\star^n c]_{i_n} \Box \dots [\star c]_{i_1} \Box [c]_{i_0} \psi$ derived using $(\text{ax-nec})^*$, we can use $(\text{ax-nec})^*$ also to obtain $[\star^{n+1} c]_i \Box \varphi$. By Lemma 1 we find $\text{LPLTL}_{\emptyset}^* \vdash \Box \varphi \rightarrow \varphi$. Hence by Lemma 24 there is a term t such that $\text{LPLTL}_{\text{CS}}^* \vdash [t]_i (\Box \varphi \rightarrow \varphi)$. We finally conclude

$$\text{LPLTL}_{\text{CS}}^* \vdash [t \cdot \star^{n+1} c]_i \varphi. \quad \square$$

It is straightforward to adapt our semantics for LPLTL_{CS} to the extended language of $\text{LPLTL}_{\text{CS}}^*$. Soundness and completeness of $\text{LPLTL}_{\text{CS}}^*$ can then be shown similar to the case of LPLTL_{CS} . However, for the completeness proof of $\text{LPLTL}_{\text{CS}}^*$ we require CS to be axiomatically appropriate in order to have the necessitation rules available.

Definition 26. Let CS be a constant specification for LPLTL^* . A CS-evidence function for agent i for LPLTL^* is a function $\mathcal{E}_i: S \times \text{Term}^* \rightarrow \mathcal{P}(\text{Fml}^*)$ satisfying conditions 1–5 of Definition 3 and the following additional condition:

- if $[c]_{i_0} \varphi \in \text{CS}$, then for all $w \in S$, all $n \geq 1$, and all agents i_{n-1}, \dots, i_1 :

$$\Box [\star^{n-1} c]_{i_{n-1}} \dots \Box [\star c]_{i_1} \Box [c]_{i_0} \varphi \in \mathcal{E}_i(w, \star^n c).$$

An $\text{LPLTL}_{\text{CS}}^*$ -interpreted system is an interpreted system where we use evidence functions for LPLTL^* . We write $\models_{\text{CS}}^* \varphi$ to mean $\mathcal{I} \models \varphi$ for all $\text{LPLTL}_{\text{CS}}^*$ -interpreted systems \mathcal{I} .

Theorem 27 (Soundness and completeness). Let CS be an axiomatically appropriate constant specification for LPLTL^* . For each formula φ ,

$$\models_{\text{CS}}^* \varphi \text{ iff } \text{LPLTL}_{\text{CS}}^* \vdash \varphi.$$

We conclude this section by showing the conservativity of LPLTL^* over LPLTL . First we need a lemma.

Lemma 28. Let CS be a constant specification for LPLTL , and \mathcal{I} be an interpreted system of LPLTL for CS. Then we can extend \mathcal{I} to an $\text{LPLTL}_{\text{CS}}^*$ -interpreted system \mathcal{I}^* such that for every run r , every $n \in \mathbb{N}$, and every formula $\varphi \in \text{Fml}$:

$$(\mathcal{I}, r, n) \models \varphi \iff (\mathcal{I}^*, r, n) \models \varphi.$$

Proof. Let $\mathcal{I} = (\mathcal{R}, S, R_1, \dots, R_h, \mathcal{E}_1, \dots, \mathcal{E}_h, \nu)$ be an arbitrary interpreted system of LPLTL for CS. By a least fixed point construction, we can easily extend the CS-evidence functions \mathcal{E}_i , for $1 \leq i \leq h$, to CS-evidence functions \mathcal{E}_i^* such that

1. $\mathcal{I}^* = (\mathcal{R}, S, R_1, \dots, R_h, \mathcal{E}_1^*, \dots, \mathcal{E}_h^*, \nu)$ is an $\text{LPLTL}_{\text{CS}}^*$ -interpreted system and
2. for each formula $\varphi \in \text{Fml}$, each run r and each $n \in \mathbb{N}$:

$$(\mathcal{I}, r, n) \models \varphi \iff (\mathcal{I}^*, r, n) \models \varphi. \quad \square$$

Theorem 29 (Conservativity). *Let CS be a constant specification for LPLTL and $\varphi \in \text{Fml}$ a formula. If $\text{LPLTL}_{CS}^* \vdash \varphi$, then $\vdash_{CS} \varphi$.*

Proof. Suppose that $\not\vdash_{CS} \varphi$. Then, by Theorem 19, we have $\not\models_{CS} \varphi$. Thus there exists an interpreted system \mathcal{J} of LPLTL for CS and a state $r(n)$ such that $(\mathcal{J}, r, n) \not\models \varphi$. Now, by Lemma 28, we find an LPLTL_{CS}^* -interpreted system \mathcal{J}^* such that $(\mathcal{J}^*, r, n) \not\models \varphi$. Therefore, by Theorem 27, we have $\text{LPLTL}_{CS}^* \not\vdash \varphi$ as desired. \square

7 Additional Principles

In LPLTL_{CS} , epistemic and temporal properties do not interact. On the other hand in LPLTL_{CS}^* , there are some interactions between time and knowledge, in axiom (boxed reflexivity) and rule (ax-nec)*. Here we propose some principles that create a connection between justifications and temporal modalities. We assume the language for terms to be augmented in the obvious way.

$$\begin{aligned}
 [t]_i \Box \varphi &\rightarrow \Box [\Downarrow t]_i \varphi && (\Box\text{-access}) \\
 \Box [t]_i \varphi &\rightarrow [\Uparrow t]_i \Box \varphi && (\text{generalize}) \\
 [t]_i \bigcirc \varphi &\rightarrow \bigcirc [\Rightarrow t]_i \varphi && (\bigcirc\text{-access}) \\
 \bigcirc [t]_i \varphi &\rightarrow [\Leftarrow t]_i \bigcirc \varphi && (\bigcirc\text{-left})
 \end{aligned}$$

Some first remarks about these principles:

- (\Box -access) This is very plausible, if you have evidence that something always is true, then at every point in time you should be able to access this information. The term operator \Downarrow makes the evidence accessible in every future point in time.
- (generalize) Using evidence this seems more plausible than just using knowledge, as one requires the evidence to be the same at every point in time. The term operator \Uparrow converts permanent evidence for a formula to evidence for believing that this formula is always true.
- (\bigcirc -access) This seems plausible: agents do not forget evidence once they have gathered it and can “take it with them”. The term operator \Rightarrow carries evidence through time.
- (\bigcirc -left) This one seems less plausible as it implies some form of premonition. The term operator \Leftarrow presages future evidence for belief.

The principle (generalize) is very strong. In particular, it makes internalization possible even in the presence of necessitation rules. Indeed, let LPLTL_{CS}^G be the system LPLTL_{CS} extended by the axioms (generalize) and (mix)—this is also reflected by constant specification—and the iterated constant necessitation rule

$$\frac{[c]_i \varphi \in CS}{[\star^n c]_{i_n} \dots [\star c]_{i_1} [c]_i \varphi}$$

for arbitrary agents i_1, \dots, i_n . Here we employ the same term operator \star as in the rule (ax-nec)* although the meaning of \star in these two rules is a bit different.

Theorem 30 (Internalization). *Let CS be an axiomatically appropriate constant specification. The system LPLTL_{CS}^G enjoys internalization.*

Proof. We proceed by induction on the derivation of φ . There are two interesting cases:

In case φ is $\Box\psi$, derived using (\Box -nec), then, by the induction hypothesis, there is a term s such that $[s]_i\psi$ is provable. Now, we can use (\Box -nec) in order to obtain $\Box[s]_i\psi$ and then (generalize) and modus ponens to get $[\uparrow s]_i\Box\psi$.

In case φ is $\bigcirc\psi$, derived using (\bigcirc -nec), then, as above, we obtain $[\uparrow s]_i\Box\psi$. Since CS is axiomatically appropriate, there is a constant c with $[c]_i(\Box\psi \rightarrow \bigcirc\psi)$. Thus we finally conclude $[c \cdot \uparrow s]_i\bigcirc\psi$. \square

It is obvious how to formulate conditions on evidence functions that correspond to the additional principles of this section such that soundness results can be obtained, see [5]. However, it is not clear how to show the existence of such models and how to show completeness for these additional principles.

8 Conclusions

We introduced the temporal justification logic LPLTL_{CS} and showed that it is sound and complete with respect to interpreted systems that are based on Fitting-models. To achieve this we had to adapt the usual canonical model construction of justification logic such that it yields a finite Fitting-model. Further, we established that a suitable form of axiom necessitation can replace the necessitation rules for \Box and \bigcirc and thus make internalization possible. Finally, we briefly discussed some additional principles that concern the interaction of knowledge, justifications, and time.

We finish this paper with some questions that show possible directions for future work.

Question 1. *How does a temporal justification logic based on JT45, i.e. the justification counterpart of the modal logic S5, look like? The problem is that JT45-models must satisfy the strong evidence condition, i.e. for all \mathcal{J}, r, n and each formula $[t]_i\varphi$*

$$\varphi \in \mathcal{E}_i(r(n), t) \text{ implies } (\mathcal{J}, r, n) \models [t]_i\varphi, \quad (7)$$

see [3, 23, 25, 26]. In infinite canonical models, the strong evidence property is an easy consequence of the Truth Lemma. In our temporal setting, we have a finite canonical model and the Truth Lemma is restricted to $\text{Sub}^+(\chi)$. Hence it does not entail (7) for all formulas $[t]_i\varphi$.

Question 2. *How can the typical examples, e.g., protocols related to message transmission, be formalized in LPLTL_{CS} ?*

Yavorskaya [27] introduces multi-agent justification logics with interaction operations on the justification terms, in particular, she studies two principles:

$$[t]_i\varphi \rightarrow \left[\uparrow_i^j t \right]_j [t]_i\varphi \quad (\text{evidence verification})$$

$$[t]_i\varphi \rightarrow \left[\uparrow_i^j t \right]_j \varphi \quad (\text{evidence conversion})$$

where one agent's evidence is converted into another agent's evidence. We believe that principles of this kind will be important in the context of this question. For example, one might consider a temporal justification logic with principles such as

$$\begin{aligned} [t]_i\varphi &\rightarrow \bigcirc [\text{sent}_j^i(t)]_j\varphi \quad \text{or} \\ [t]_i\varphi &\rightarrow \Diamond [\text{sent}_j^i(t)]_j\varphi. \end{aligned}$$

Here agent i sends evidence t for φ to agent j and the term $\text{sent}_j^i(t)$ denotes the evidence that agent j received for believing φ .

Question 3. *What happens if we require operations on justification terms to take time?*

We could formalize this idea, e.g., by replacing (application), (sum), and (positive introspection) with

$$\begin{aligned} [t]_i(\varphi \rightarrow \psi) &\rightarrow ([s]_i\varphi \rightarrow \bigcirc [t \cdot s]_i\psi) \\ [t]_i\varphi \vee [s]_i\varphi &\rightarrow \bigcirc [t + s]_i\varphi \\ [t]_i\varphi &\rightarrow \bigcirc [!t]_i[t]_i\varphi. \end{aligned}$$

This might also relate to the logical omniscience problem [4].

Question 4. *Can dynamic epistemic justification logics be translated into temporal justification logic akin to [9]?*

There are several dynamic justification logics available, e.g., [7, 8, 20, 24], which feature not only traditional public announcements but also specific forms of evidence based updates and evidence elimination. It would be interesting to see what the relationship between those dynamic logics and temporal justification logic is.

This paper showed a first successful combination of temporal and justification logic. While this initial work shows the feasibility of combining these logics with minimal interaction, the list of questions above shows that various interesting properties may arise from more intricate interactions between justified knowledge and time.

References

- [1] Sergei N. Artemov (2001): *Explicit Provability and Constructive Semantics*. *Bulletin of Symbolic Logic* 7(1), pp. 1–36.
- [2] Sergei [N.] Artemov (2006): *Justified common knowledge*. *Theoretical Computer Science* 357(1–3), pp. 4–22, doi:10.1016/j.tcs.2006.03.009.
- [3] Sergei [N.] Artemov (2008): *The Logic of Justification*. *The Review of Symbolic Logic* 1(4), pp. 477–513, doi:10.1017/S1755020308090060.
- [4] Sergei [N.] Artemov & Roman Kuznets (2014): *Logical omniscience as infeasibility*. *Annals of Pure and Applied Logic* 165(1), pp. 6–25, doi:10.1016/j.apal.2013.07.003. Published online August 2013.
- [5] Samuel Bucheli (2015): *Some Notes on Temporal Justification Logic*. CoRR abs/1510.07247. Available at <http://arxiv.org/abs/1510.07247>.
- [6] Samuel Bucheli, Roman Kuznets & Thomas Studer (2011): *Justifications for common knowledge*. *Journal of Applied Non-Classical Logics* 21(1), pp. 35–60, doi:10.3166/JANCL.21.35-60.
- [7] Samuel Bucheli, Roman Kuznets & Thomas Studer (2011): *Partial Realization in Dynamic Justification Logic*. In Lev D. Beklemishev & Ruy de Queiroz, editors: *WoLLIC 2011, Proceedings, LNAI 6642*, Springer, pp. 35–51, doi:10.1007/978-3-642-20920-8_9.
- [8] Samuel Bucheli, Roman Kuznets & Thomas Studer (2014): *Realizing public announcements by justifications*. *Journal of Computer and System Sciences* 80(6), pp. 1046–1066, doi:http://dx.doi.org/10.1016/j.jcss.2014.04.001.
- [9] Hans van Ditmarsch, Wiebe van der Hoek & Ji Ruan (2013): *Connecting dynamic epistemic and temporal epistemic logics*. *Logic Journal of IGPL* 21(3), pp. 380–403, doi:10.1093/jigpal/jzr038.
- [10] Ronald Fagin, Joseph Y. Halpern, Yoram Moses & Moshe Y. Vardi (1995): *Reasoning about Knowledge*. MIT Press.
- [11] Melvin Fitting (2005): *The logic of proofs, semantically*. *Annals of Pure and Applied Logic* 132(1), pp. 1–25, doi:10.1016/j.apal.2004.04.009.

- [12] Dov M. Gabbay, Ian Hodkinson & Mark Reynolds (1994): *Temporal Logic (Vol. 1): Mathematical Foundations and Computational Aspects*. Oxford University Press, Inc.
- [13] Meghdad Ghari (2014): *Distributed Knowledge Justification Logics*. *Theory of Computing Systems* 55(1), pp. 1–40, doi:10.1007/s00224-013-9492-x.
- [14] Meghdad Ghari (2016): *Pavelka-style fuzzy justification logics*. *Logic Journal of IGPL* 24(5), pp. 743–773, doi:10.1093/jigpal/jzw019.
- [15] Robert Goldblatt (1992): *Logics of Time and Computation*, 2nd edition. Center for the Study of Language and Information.
- [16] Rajeev Goré (1999): *Tableau Methods for Modal and Temporal Logics*. In Marcello D’Agostino, Dov M. Gabbay, Reiner Hähnle & Joachim Posegga, editors: *Handbook of Tableau Methods*, Springer Netherlands, pp. 297–396, doi:10.1007/978-94-017-1754-0_6.
- [17] Joseph Y. Halpern, Ron van der Meyden & Moshe Y. Vardi (2004): *Complete Axiomatizations for Reasoning About Knowledge and Time*. *SIAM J. Comput.* 33(3), pp. 674–703, doi:10.1137/S0097539797320906.
- [18] Joseph Y. Halpern & Lenore D. Zuck (1992): *A Little Knowledge Goes a Long Way: Knowledge-based Derivations and Correctness Proofs for a Family of Protocols*. *J. ACM* 39(3), pp. 449–478, doi:10.1145/146637.146638.
- [19] Ioannis Kokkinis, Petar Maksimović, Zoran Ognjanović & Thomas Studer (2015): *First steps towards probabilistic justification logic*. *Logic Journal of IGPL* 23(4), pp. 662–687, doi:10.1093/jigpal/jzv025.
- [20] Roman Kuznets & Thomas Studer (2013): *Update as Evidence: Belief Expansion*. In Sergei [N.] Artemov & Anil Nerode, editors: *LFCS 2013, Proceedings, LNCS 7734*, Springer, pp. 266–279, doi:10.1007/978-3-642-35722-0_19.
- [21] Ron van der Meyden & Ka-shu Wong (2003): *Complete Axiomatizations for Reasoning about Knowledge and Branching Time*. *Studia Logica* 75(1), pp. 93–123.
- [22] Alexey Mkrtychev (1997): *Models for the Logic of Proofs*. In Sergei Adian & Anil Nerode, editors: *LFCS’97, Proceedings, LNCS 1234*, Springer, pp. 266–275, doi:10.1007/3-540-63045-7_27.
- [23] Eric Pacuit (2005): *A Note on Some Explicit Modal Logics*. In: *Proceedings of the 5th Panhellenic Logic Symposium*, University of Athens, Athens, Greece, pp. 117–125.
- [24] Bryan Renne (2012): *Multi-agent Justification Logic: communication and evidence elimination*. *Synthese* 185(S1), pp. 43–82, doi:10.1007/s11229-011-9968-7. Published online July 2011.
- [25] Natalia [M.] Rubtsova (2006): *On Realization of S5-modality by Evidence Terms*. *Journal of Logic and Computation* 16(5), pp. 671–684, doi:10.1093/logcom/exl030.
- [26] Thomas Studer (2013): *Decidability for Some Justification Logics with Negative Introspection*. *Journal of Symbolic Logic* 78(2), pp. 388–402, doi:10.2178/jsl.7802030.
- [27] Tatiana Yavorskaya (Sidon) (2008): *Interacting Explicit Evidence Systems*. *Theory of Computing Systems* 43(2), pp. 272–293, doi:10.1007/s00224-007-9057-y. Published online October 2007.